

## Literatur

zum Beitrag von Dr. E. Blaurock-Busch „GADOLINIUM-Kontrastmittel – Ablagerung und Entgiftung“

<https://www.dkfz.de/de/radiologie/>

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/019596s063lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/019596s063lbl.pdf)

Eliasz I, Hotchkiss AT, Fishman ML, Rode D. The effect of modified citrus pectin on urinary excretion of toxic elements. *Phytoth Res.* 2006; 20:859-864

Dawson P, Punwani Sh. Nephrogenic systemic fibrosis: non-gadolinium options for the imaging of CKD/ESRD patients. *Semin Dial* 2008 Arch-Apr: 21(2):160-5

Khotimchenko Y, Khozhaenko E, Kovalev V, Khotimchenko M. Cerium Binding Activity of Pectins Isolated from the Seagrasses *Zostera marina* and *Phyllospadix iwatensis*. *Mar Drugs* 2012; 10:834-848

Kusrine E, Wicaksone W, Gunawan C et al. Kinetics, mechanism, and thermodynamics of lanthanum adsorption on pectin extracted from durian rind. *J Environm Chem Engineering* 2018 (6):6580-6588

Murata N et al. (2016). Macrocyclic and Other Non-Group 1 Gadolinium Contrast Agents Deposit Low Levels of Gadolinium in Brain and Bone Tissue: Preliminary Results From 9 Patients With Normal Renal Function. *Invest Radiol.* Feb 8, epub ahead of print